

学校编码：10384

学号：32420131152302

廈門大學

硕士学位论文

淀粉和葡萄糖化学转化为果糖的研究

Study on the Conversion of Starch and Glucose into Fructose

陈晖晖

指导教师：林鹿

专业名称：工程硕士(材料工程)

答辩日期：2016年5月

厦门大学学位论文原创性声明

本人呈交的学位论文是本人在导师指导下，独立完成的研究成果。本人在论文写作中参考其他个人或集体已经发表的研究成果，均在文中以适当方式明确标明，并符合法律规范和《厦门大学研究生学术活动规范(试行)》。

另外，该学位论文为()课题(组)的研究成果，获得()课题(组)经费或实验室的资助，在()实验室完成。(请在以上括号内填写课题或课题组负责人或实验室名称，未有此项声明内容的，可以不作特别声明。)

声明人(签名)：

年 月 日

厦门大学学位论文著作权使用声明

本人同意厦门大学根据《中华人民共和国学位条例暂行实施办法》等规定保留和使用此学位论文，并向主管部门或其指定机构送交学位论文(包括纸质版和电子版)，允许学位论文进入厦门大学图书馆及其数据库被查阅、借阅。本人同意厦门大学将学位论文加入全国博士、硕士学位论文共建单位数据库进行检索，将学位论文的标题和摘要汇编出版，采用影印、缩印或者其它方式合理复制学位论文。

本学位论文属于：

()1. 经厦门大学保密委员会审查核定的保密学位论文，于
年 月 日解密，解密后适用上述授权。

()2. 不保密，适用上述授权。

(请在以上相应括号内打“√”或填上相应内容。保密学位论文应是已经厦门大学保密委员会审定过的学位论文，未经厦门大学保密委员会审定的学位论文均为公开学位论文。此声明栏不填写的，默认为公开学位论文，均适用上述授权。)

声明人(签名)：

年 月 日

摘 要

在全球经济和工业高速发展的同时，化石能源日渐减少，环境污染和生态恶化等问题日益严重，因此寻找可再生资源以缓解当前的能源和环境危机迫在眉睫。生物质资源被认为唯一能替代化石能源以获取高附加值化学品、生物燃料和能源材料的绿色、可再生资源。碳水化合物是生物质资源中储量最丰富的组分，而淀粉是人类饮食中最为丰富的碳水化合物，将其转化为果糖是当前开发利用生物质资源的研究热点。果糖作为生物质重要的衍生化学品之一，是一种重要的食品甜味剂和制备生物基化学品最为活跃的原材料，市场上对果糖的需求不断扩大，其在燃料和能源材料等方面的应用也逐渐被挖掘出来。碳水化合物先水解为葡萄糖而后葡萄糖通过异构化反应转化为果糖，葡萄糖异构化为果糖是碳水化合物转化为平台化学品和生物燃料关键的中间步骤。本论文以绿色化学为宗旨，以淀粉和葡萄糖为原料，针对目前葡萄糖异构化反应中存在的问题展开了一系列研究，包括绿色环保、价格低廉和制备简单的催化剂的开发，各步反应条件的优化以及5-羟甲基糠醛（HMF）制备体系的建立等，并取得了一定的研究成果。

首先，在水体系中，以硫酸湿法浸渍沸石分子筛得到一类新型负载催化剂，能催化淀粉一锅法制备果糖，发现固体酸催化剂S042-/USY表现出良好的催化活性，果糖的得率最高。各种反应条件对果糖的得率都具有较大的影响，在5 wt.%淀粉（淀粉/水）、催化剂用量为淀粉质量的30 wt.%、反应温度为150 °C、反应时间为1 h、转速为400 rpm的条件下，能得到得率最高的果葡糖浆，含27.84%的果糖和58.34%的葡萄糖。通过对催化剂进行物理吸附表征，发现USY具有较高的孔隙度和比表面积，通过浸渍焙烧过程能有效使S042-与USY通过键合作用吸附在USY表面上。同时催化剂在使用过程中表现出重复使用性能不佳的缺点，通过对催化剂进行化学吸附和元素分析表征，发现催化剂活性的降低与含碳有机质沉积和S042-催化活性中心的流失有关。在此基础上开发了一套通过简单的焙烧（除去表面积碳）和硫酸中浸渍活化（增加S042-）的催化剂再生方法，可以有效地恢复S042-/USY的催化活性。

其次，建立了淀粉两步法制备果糖的高效反应路线。第一步是用超低酸水解淀粉制

备葡萄糖，第二步是用碱性催化剂催化葡萄糖异构化为果糖。第一步当淀粉浓度为5 wt. %，硫酸浓度为8 mM，在150 °C反应1 h，能获得103.83%的葡萄糖得率，淀粉几乎完全水解为葡萄糖。第二步选取实验室常见的碱性催化剂作为考察对象，发现一系列强碱弱酸盐和有机碱对葡萄糖异构化反应均具有良好的催化效果。在强碱弱酸盐中选用安全无毒并且易于去除的碳酸钠为催化剂，当葡萄糖浓度为5 wt. %（葡萄糖/水），碳酸钠用量为35 wt. %（碳酸钠/葡萄糖），在65 °C反应3 h，葡萄糖转化率为62.37%，果糖选择性为64.34%，果糖得率为40.13%。在有机碱中选用可作为营养品添加剂的胆碱为催化剂，当葡萄糖浓度为5 wt. %（葡萄糖/水），胆碱用量为10 wt. %（胆碱/葡萄糖），在65 °C反应3 h，葡萄糖转化率为53.25%，果糖选择性为80.38%，果糖得率为42.8%。同时，研究发现：葡萄糖的异构化反应与催化剂的种类无关，而与反应体系pH值有很大相关性。只要在合适的pH值范围就可以得到高得率的果糖。

然后，在以胆碱为催化剂的基础上添加一定量的硼砂能够进一步提高葡萄糖异构化效率，证明硼砂具有良好的助催化作用。另外，根据硼砂的特性，提出了硼砂助催化葡萄糖异构化制备果糖的可能反应机理。

最后，在以上实验研究的基础上，建立了葡萄糖两步制备HMF的温和、高效的反应路线。第一步用胆碱和硼砂协同催化葡萄糖异构化为果糖；第二步将生成的果糖与氯化胆碱形成低共熔溶剂制备HMF。其中，在氯化胆碱4倍量wt/wt（氯化胆碱/生成的果糖）、1 wt. %盐酸（盐酸/果糖）、100 °C反应4 h的条件下，能获得70.8 mol%的HMF得率。胆碱和氯化胆碱催化体系与目前报道的氯化铬离子液体催化体系相比，避免了有毒铬离子催化剂污染环境的问题且具有方法简单、条件温和、价格低廉、绿色环保等优点，为今后HMF的规模化生产提供了一定的理论指导和可行措施。

关键词：淀粉；固体酸；葡萄糖；异构化；果糖；碱；硼砂；p；HMF

Abstract

With the fast development of world economy, the diminishment of fossil resources and the gradual deterioration of global ecological environment, finding renewable resources to alleviate the current energy and environmental crisis is imminent. Biomass is considered to be the only renewable resources in nature to replace the fossil fuels for the production of high value added chemicals, biofuels and energy materials. The conversion of carbohydrates, which are the most abundant components of biomass, into fructose is one of the most important ways for the utilization of biomass energy. As fructose is an important derivative of biomass, an important sweetener and active raw material for the production of biologic-chemistry matter, its application in food, fuels and energy materials has been gradually excavated. Starch, which is the most abundant component of carbohydrates, can be hydrolyzed to glucose and glucose can be converted to fructose by isomerization. The isomerization of glucose into fructose is considered to be a critical intermediate step in the efficient conversion of renewable lignocellulosic materials into biofuels and platform chemicals. According to the current research progress and existing problems in the isomerization of glucose into fructose, a series of studies on the concept of green chemistry have been carried out in this dissertation. Various low-budget, environmentally friendly, easily prepared catalysts were prepared for the conversion of starch and glucose into fructose, the basic reaction conditions of every step were optimized and the reaction system for the preparation of HMF was established. Firstly, supporting sulfuric acid on zeolite was found to be an efficient catalyst to bring about the conversion of starch directly into fructose in a one-step reaction. The catalytic efficiency of a series of modified zeolite catalysts for the conversion of starch into fructose was investigated in details. Among these catalysts used, the SO42-/USY catalyst possessed a bifunction with high activity for both

hydrolysis of starch and isomerization of glucose, which enable one-step preparation of HFCS from starch to turn into reality. An optimized fructose yield of 27.84% was obtained from starch under the following reaction conditions: 5 wt.% starch in water, 30 wt.% SO₄²⁻/USY (SO₄²⁻/USY /starch), 150 °C, 1 h, 400 rpm. The USY catalyst was found has high specific surface area and porosity characterized by physical adsorption technique, which enable the sulfuric acid to be effectively adsorbed on the USY surface by means of impregnation and calcination. Meanwhile, the catalysts were characterized by NH₃-TPD analysis and elemental analysis to find that the decrease of catalytic activity resulted from carbon deposition on the catalysts and the loss of SO₄²⁻. Therefore, the catalytic activity of SO₄²⁻/USY can be effectively recovered by calcining (removal carbon on the surface of catalyst) and impregnating in sulfuric acid (increase SO₄²⁻).

Secondly, a high efficient reaction route for the two-step preparation of fructose from starch was established. Step 1, the starch was hydrolyzed into glucose by ultra low acid; step 2, the glucose was isomerized to fructose by basic catalyst. In the first step, the glucose yield of 103.83% (starch was almost completely hydrolyzed to glucose) was obtained under the optimal reaction conditions: 5 wt.% starch in water, 8mM sulfuric acid concentration, 150 °C, 1 h. In the second step, a series of alkali salt and organic alkali were found to have good catalytic efficiency for the isomerization of glucose. Among alkali salt, sodium carbonate was selected as the best catalyst because of the obvious advantages of safety, innocuity and easy to removal. A glucose conversion of 62.37%, a fructose selectivity of 64.34% and a fructose yield of 40.13% were obtained under the following reaction conditions: 5 wt.% glucose in water, 35 wt.% Na₂CO₃ (Na₂CO₃/glucose), 65 °C, 3 h. Among organic alkali, choline was chosen as the optimal catalyst because it can be used as a nutritional additive. A glucose conversion of 53.25%, a fructose selectivity of 80.38% and a fructose yield of 42.8% were obtained under the following reaction conditions: 5 wt.% glucose in

water, 10 wt.% choline (choline/glucose), 65 °C, 3 h. Meanwhile, the study found that with proper pH value, almost all of inorganic bases, organic bases and alkali salt could efficiently isomerize glucose into fructose. The results indicated that the isomerization process was significantly affected by the basicity of reaction system, while the types of alkalines had little effect.

Subsequently, borax could enhance the catalytic efficiency of glucose isomerization by choline. According to the characteristics of borax, the possible reaction mechanism for borax promoting effects on the glucose isomerization was presented.

Eventually, according to the study aforementioned, a mild and high efficient reaction route for the two-step preparation of HMF from glucose was designed and implemented. Step 1, the glucose was isomerized to fructose by choline and borax; step 2, the fructose was converted to HMF with hydrochloric acid induced by choline chloride. An optimized HMF yield of 70.8 mol% was obtained from glucose under the following reaction conditions: 4 wt/wt ChCl (ChCl/fructose), 1 wt.% HCl (HCl/fructose), 100 °C, 4 h. Compared with the preparation of HMF from glucose in metal chloride/ionic liquid reaction system, this reaction system possesses the characteristics of easy process, low investment and benign to environment. This study was drafted to provide a theoretical basis and technical support for the mass production of HMF in the future.

Keywords: starch; solid acid; glucose; isomerization; fructose; base; choline; pH; HMF

参考资料

- [1] G. Fischer, L. Schrattenholzer, Global bioenergy potentials through 2050, *Biomass and bioenergy*, 20 (2001) 151-159.
- [2] Y. Román-Leshkov, C.J. Barrett, Z.Y. Liu, J.A. Dumesic, Production of dimethylfuran for liquid fuels from biomass-derived carbohydrates, *Nature*, 447 (2007) 982-985.
- [3] C.B. Field, J.E. Campbell, D.B. Lobell, Biomass energy: the scale of the potential resource, *Trends in ecology & evolution*, 23 (2008) 65-72.
- [4] S. Liu, Woody biomass conversion: sustainability and water-based processes, *Journal of Bioprocess Engineering and Biorefinery*, 1 (2012) 6-32.
- [5] J.S. Yuan, K.H. Tiller, H. Al-Ahmad, N.R. Stewart, C.N. Stewart Jr, Plants to power: bioenergy to fuel the future, *Trends in plant science*, 13 (2008) 421-429.
- [6] A.J. Ragauskas, C.K. Williams, B.H. Davison, G. Britovsek, J. Cairney, C.A. Eckert, W.J. Frederick, J.P. Hallett, D.J. Leak, C.L. Liotta, The path forward for biofuels and biomaterials, *science*, 311 (2006) 484-489.
- [7] J.B. Binder, R.T. Raines, Simple chemical transformation of lignocellulosic biomass into furans for fuels and chemicals, *Journal of the American Chemical Society*, 131 (2009) 1979-1985.
- [8] S. Liu, H. Lu, R. Hu, A. Shupe, L. Lin, B. Liang, A sustainable woody biomass biorefinery, *Biotechnology Advances*, 30 (2012) 785-810.
- [9] J.Y.G. Chan, Y. Zhang, Selective Conversion of Fructose to 5-Hydroxymethylfurfural Catalyzed by Tungsten Salts at Low Temperatures, *ChemSusChem*, 2 (2009) 731-734.
- [10] H. Amiri, K. Karimi, S. Roodpeyma, Production of furans from rice straw by single-phase and biphasic systems, *Carbohydrate research*, 345 (2010) 2133-2138.
- [11] F. Tao, H. Song, L. Chou, Catalytic conversion of cellulose to chemicals in ionic liquid, *Carbohydrate research*, 346 (2011) 58-63.
- [12] H.L. Chum, R.P. Overend, Biomass and renewable fuels, *Fuel processing technology*, 71 (2001) 187-195.
- [13] A. Demirbağ, Biomass resource facilities and biomass conversion processing for fuels and chemicals, *Energy conversion and management*, 42 (2001) 1357-1378.
- [14] 张燕平, 变性淀粉制造与应用, 北京: 化学工业出版社, 2001.
- [15] R. Whistler, 淀粉的化学与工艺学, 北京: 中国食品出版社, 1988.
- [16] T.P. Binder, A. Sanborn, Processes for making sugar and/or sugar alcohol dehydration products, in, Google Patents, 2013.
- [17] J. Straatsma, K. Vellenga, H. De Wilt, G. Joosten, Isomerization of glucose to fructose. 2. Optimization of reaction conditions in the production of high fructose syrup by isomerization of glucose catalyzed by a whole cell immobilized glucose isomerase catalyst, *Industrial & Engineering Chemistry Process Design and Development*, 22 (1983) 356-361.
- [18] M.G. Tordoff, A.M. Alleva, Effect of drinking soda sweetened with aspartame or high-fructose corn syrup on food intake and body weight, *The American Journal of Clinical Nutrition*, 51 (1990) 963-969.
- [19] E. Nikolla, Y. Román-Leshkov, M. Moliner, M.E. Davis, "One-pot" synthesis of 5-(hydroxymethyl) furfural from carbohydrates using tin-beta zeolite, *Acs Catalysis*, 1 (2011) 408-410.
- [20] Z.-p. GAO, T.-l. YUE, Y.-h. YUAN, Y.-y. WANG, Research progress on producing technology and the applied characteristics of fructose [J], *Journal of Northwest Sci-Tech University of Agriculture and Forestry S*, 1 (2003).
- [21] 赵锡武, 何玉莲, 果糖的特性及应用, *饮料工业*, 9 (2006) 4-9.
- [22] A. Corma, S. Iborra, A. Velty, Chemical routes for the transformation of biomass into chemicals, *Chemical Reviews*, 107 (2007) 2411-2502.
- [23] A. Takagaki, M. Ohara, S. Nishimura, K. Ebitani, A one-pot reaction for biorefinery: combination of solid acid and base catalysts for direct production of 5-hydroxymethylfurfural from saccharides, *Chem. Commun.*,

(2009) 6276-6278.

- [24] A. Takagaki, M. Takahashi, S. Nishimura, K. Ebitani, One-pot synthesis of 2, 5-diformylfuran from carbohydrate derivatives by sulfonated resin and hydrotalcite-supported ruthenium catalysts, *ACS Catalysis*, 1 (2011) 1562-1565.
- [25] Q. Ren, Y. Huang, H. Ma, F. Wang, J. Gao, J. Xu, Conversion of glucose to 5-hydroxymethylfurfural catalyzed by metal halide in N, N-dimethylacetamide, *BioResources*, 8 (2013) 1563-1572.
- [26] H. Mehdi, V. F á bos, R. Tuba, A. Bodor, L.T. Mika, I.T. Horv á th, Integration of homogeneous and heterogeneous catalytic processes for a multi-step conversion of biomass: From sucrose to levulinic acid, - valerolactone, 1, 4-pentanediol, 2-methyl-tetrahydrofuran, and alkanes, *Topics in Catalysis*, 48 (2008) 49-54.
- [27] J. Zhang, S. Wu, B. Li, H. Zhang, Advances in the catalytic production of valuable levulinic acid derivatives, *ChemCatChem*, 4 (2012) 1230-1237.
- [28] P.A. Son, S. Nishimura, K. Ebitani, Synthesis of levulinic acid from fructose using Amberlyst-15 as a solid acid catalyst, *Reaction Kinetics, Mechanisms and Catalysis*, 106 (2012) 185-192.
- [29] F. Guo, Z. Fang, T.-J. Zhou, Conversion of fructose and glucose into 5-hydroxymethylfurfural with lignin-derived carbonaceous catalyst under microwave irradiation in dimethyl sulfoxide – ionic liquid mixtures, *Bioresource technology*, 112 (2012) 313-318.
- [30] L. Hu, Y. Sun, L. Lin, S. Liu, 12-Tungstophosphoric acid/boric acid as synergetic catalysts for the conversion of glucose into 5-hydroxymethylfurfural in ionic liquid, biomass and bioenergy, 47 (2012) 289-294.
- [31] A.S. Amarasekara, L.D. Williams, C.C. Ebede, Mechanism of the dehydration of d-fructose to 5-hydroxymethylfurfural in dimethyl sulfoxide at 150°C: an NMR study, *Carbohydrate research*, 343 (2008) 3021-3024.
- [32] M.J. Antal, W.S. Mok, G.N. Richards, Mechanism of formation of 5-(hydroxymethyl)-2-furaldehyde from D-fructose and sucrose, *Carbohydrate research*, 199 (1990) 91-109.
- [33] R.S. Assary, P.C. Redfern, J. Greeley, L.A. Curtiss, Mechanistic insights into the decomposition of fructose to hydroxy methyl furfural in neutral and acidic environments using high-level quantum chemical methods, *The Journal of Physical Chemistry B*, 115 (2011) 4341-4349.
- [34] J. Guan, Q. Cao, X. Guo, X. Mu, The mechanism of glucose conversion to 5-hydroxymethylfurfural catalyzed by metal chlorides in ionic liquid: A theoretical study, *Computational and Theoretical Chemistry*, 963 (2011) 453-462.
- [35] X. Qian, Mechanisms and Energetics for Brønsted Acid-Catalyzed Glucose Condensation, Dehydration and Isomerization Reactions, *Topics in Catalysis*, 55 (2012) 218-226.
- [36] C. Moreau, R. Durand, S. Razigade, J. Duhamet, P. Faugeras, P. Rivalier, P. Ros, G. Avignon, Dehydration of fructose to 5-hydroxymethylfurfural over H-mordenites, *Applied Catalysis A: General*, 145 (1996) 211-224.
- [37] S. Hu, Z. Zhang, J. Song, Y. Zhou, B. Han, Efficient conversion of glucose into 5-hydroxymethylfurfural catalyzed by a common Lewis acid SnCl₄ in an ionic liquid, *Green Chemistry*, 11 (2009) 1746-1749.
- [38] X. Qi, M. Watanabe, T.M. Aida, R.L. Smith, Fast Transformation of Glucose and Di /Polysaccharides into 5 Hydroxymethylfurfural by Microwave Heating in an Ionic Liquid/Catalyst System, *ChemSusChem*, 3 (2010) 1071-1077.
- [39] H. Zhao, J.E. Holladay, H. Brown, Z.C. Zhang, Metal chlorides in ionic liquid solvents convert sugars to 5-hydroxymethylfurfural, *Science*, 316 (2007) 1597-1600.
- [40] T. Ståhlberg, S. Rodriguez Rodriguez, P. Fristrup, A. Riisager, Metal Free Dehydration of Glucose to 5 (Hydroxymethyl) furfural in Ionic Liquids with Boric Acid as a Promoter, *Chemistry-A European Journal*, 17 (2011) 1456-1464.
- [41] Y. Rom á n Leshkov, M. Moliner, J.A. Labinger, M.E. Davis, Mechanism of glucose isomerization using a solid Lewis acid catalyst in water, *Angewandte Chemie International Edition*, 49 (2010) 8954-8957.
- [42] E.A. Pidko, V. Degirmenci, R.A. van Santen, E.J. Hensen, Glucose activation by transient Cr²⁺ dimers, *Angewandte Chemie International Edition*, 49 (2010) 2530-2534.
- [43] M. Moliner, Y. Rom á n-Leshkov, M.E. Davis, Tin-containing zeolites are highly active catalysts for the

- isomerization of glucose in water, *Proceedings of the National Academy of Sciences*, 107 (2010) 6164-6168.
- [44] J.B. Binder, A.V. Cefali, J.J. Blank, R.T. Raines, Mechanistic insights on the conversion of sugars into 5-hydroxymethylfurfural, *Energy & Environmental Science*, 3 (2010) 765-771.
- [45] K. Buchholz, J. Seibel, Industrial carbohydrate biotransformations, *Carbohydrate research*, 343 (2008) 1966-1979.
- [46] Y. Zhang, K. Hidajat, A.K. Ray, Optimal design and operation of SMB bioreactor: production of high fructose syrup by isomerization of glucose, *Biochemical engineering journal*, 21 (2004) 111-121.
- [47] P. Biely, In JR Whitaker, AGJ Voragen, & DWS Wong (Eds.) *Handbook of food enzymology* pp. 879 – 915, New York: Marcel Dekker, 2003.
- [48] B.M. Kabyemela, T. Adschiri, R.M. Malaluan, K. Arai, Glucose and fructose decomposition in subcritical and supercritical water: detailed reaction pathway, mechanisms, and kinetics, *Industrial & Engineering Chemistry Research*, 38 (1999) 2888-2895.
- [49] J.C. Sowden, R. Schaffer, The Isomerization of D-Glucose by Alkali in D₂O at 25°C, *Journal of the American Chemical Society*, 74 (1952) 505-507.
- [50] B.Y. Yang, R. Montgomery, Alkaline degradation of fructofuranosides, *Carbohydrate research*, 280 (1996) 47-57.
- [51] Y. Kanie, K. Akiyama, M. Iwamoto, Reaction pathways of glucose and fructose on Pt nanoparticles in subcritical water under a hydrogen atmosphere, *Catalysis Today*, 178 (2011) 58-63.
- [52] R. Nagorski, J.P. Richard, Mechanistic imperatives for aldose-ketose isomerization in water: Specific, general base-and metal ion-catalyzed isomerization of glyceraldehyde with proton and hydride transfer, *Journal of the American Chemical Society*, 123 (2001) 794-802.
- [53] G. Akiyama, R. Matsuda, H. Sato, S. Kitagawa, Catalytic Glucose Isomerization by Porous Coordination Polymers with Open Metal Sites, *Chemistry – An Asian Journal*, 9 (2014) 2772-2777.
- [54] D.W. Harris, M.S. Feather, Evidence for a C-2 → C-1 intramolecular hydrogen-transfer during the acid-catalyzed isomerization of D-glucose to D-fructose, *Carbohydrate Research*, 30 (1973) 359-365.
- [55] C. de Bruyn, W.A. Van Ekenstein, Action des alcalis sur les sucres, II. Transformation réciproque des uns dans les autres des sucres glucose, fructose et mannose, *Recueil des Travaux Chimiques des Pays-Bas*, 14 (1895) 203-216.
- [56] A. Onda, T. Ochi, K. Kajiyoshi, K. Yanagisawa, A new chemical process for catalytic conversion of D-glucose into lactic acid and gluconic acid, *Applied catalysis A: general*, 343 (2008) 49-54.
- [57] Isomerization of glucose, maltose, and lactose with amino compounds, *Google Patents*, 1970.
- [58] C. Moreau, R. Durand, A. Roux, D. Tichit, Isomerization of glucose into fructose in the presence of cation-exchanged zeolites and hydrotalcites, *Applied Catalysis A: General*, 193 (2000) 257-264.
- [59] J. Beenackers, B. Kuster, H. van der Baan, Physical properties of anion exchangers used as a catalyst in the isomerization of hexoses, *Applied catalysis*, 16 (1985) 75-87.
- [60] A. condensations over reconstructed Mg-Al, hydrotalcites: structure-activity relationships related to the rehydration method Abello, Sonia.
- [61] S. Yu, E. Kim, S. Park, I.K. Song, J.C. Jung, Isomerization of glucose into fructose over Mg – Al hydrotalcite catalysts, *Catalysis Communications*, 29 (2012) 63-67.
- [62] J. Di Cosimo, V. D'Amico, M. Xu, E. Iglesia, C. Apesteguía, Structure and surface and catalytic properties of Mg-Al basic oxides, *Journal of Catalysis*, 178 (1998) 499-510.
- [63] S. Saravanamurugan, M. Paniagua, J.A. Melero, A. Riisager, Efficient isomerization of glucose to fructose over zeolites in consecutive reactions in alcohol and aqueous media, *Journal of the American Chemical Society*, 135 (2013) 5246-5249.
- [64] P. Kunkeler, B. Zuurdeeg, J. Van Der Waal, J. Van Bokhoven, D. Koningsberger, H. Van Bekkum, Zeolite beta: the relationship between calcination procedure, aluminum configuration, and Lewis acidity, *Journal of catalysis*, 180 (1998) 234-244.
- [65] A. Corma, M.E. Domine, L. Nemeth, S. Valencia, Al-free Sn-Beta zeolite as a catalyst for the selective

- reduction of carbonyl compounds (Meerwein-Ponndorf-Verley reaction), *Journal of the American Chemical Society*, 124 (2002) 3194-3195.
- [66] J. Lecomte, A. Finiels, C. Moreau, Kinetic Study of the Isomerization of Glucose into Fructose in the Presence of Anion modified Hydrotalcites, *Starch*, 54 (2002) 75-79.
- [67] A.M.G. Pedrosa, M.J. Souza, D.M. Melo, A.S. Araujo, Cobalt and nickel supported on HY zeolite: Synthesis, characterization and catalytic properties, *Materials research bulletin*, 41 (2006) 1105-1111.
- [68] J. Pérez-Ramírez, S. Abelló, N.M. van der Pers, Influence of the divalent cation on the thermal activation and reconstruction of hydrotalcite-like compounds, *The Journal of Physical Chemistry C*, 111 (2007) 3642-3650.
- [69] N. Rai, S. Caratzoulas, D.G. Vlachos, Role of silanol group in Sn-beta zeolite for glucose isomerization and epimerization reactions, *ACS Catalysis*, 3 (2013) 2294-2298.
- [70] G. Li, E.A. Pidko, E.J. Hensen, Synergy between Lewis acid sites and hydroxyl groups for the isomerization of glucose to fructose over Sn-containing zeolites: a theoretical perspective, *Catalysis Science & Technology*, 4 (2014) 2241-2250.
- [71] M. Liu, S. Jia, C. Li, A. Zhang, C. Song, X. Guo, Facile preparation of Sn-zeolites by post-synthesis (isomorphous substitution) method for isomerization of glucose to fructose, *Chinese Journal of Catalysis*, 35 (2014) 723-732.
- [72] J. Dijkmans, M. Dusselier, D. Gabriëls, K. Houthoofd, P.C. Magusin, S. Huang, Y. Pontikes, M. Trekels, A. Vantomme, L. Giebler, Cooperative Catalysis for Multistep Biomass Conversion with Sn/Al Beta Zeolite, *ACS Catalysis*, 5 (2015) 928-940.
- [73] T. Ennaert, J. Geboers, E. Gobechiya, C.M. Courtin, M. Kurttepli, K. Houthoofd, C.E. Kirschhock, P.C. Magusin, S. Bals, P.A. Jacobs, Conceptual Frame Rationalizing the Self-Stabilization of H-USY Zeolites in Hot Liquid Water, *ACS Catalysis*, 5 (2014) 754-768.
- [74] M. Watanabe, Y. Aizawa, T. Iida, R. Nishimura, H. Inomata, Catalytic glucose and fructose conversions with TiO₂ and ZrO₂ in water at 473K: Relationship between reactivity and acid – base property determined by TPD measurement, *Applied Catalysis A: General*, 295 (2005) 150-156.
- [75] A. Chareonlimkun, V. Champreda, A. Shotipruk, N. Laosiripojana, Catalytic conversion of sugarcane bagasse, rice husk and corncob in the presence of TiO₂, ZrO₂ and mixed-oxide TiO₂ – ZrO₂ under hot compressed water (HCW) condition, *Bioresource technology*, 101 (2010) 4179-4186.
- [76] X. Qi, M. Watanabe, T.M. Aida, R.L. Smith, Catalytic conversion of fructose and glucose into 5-hydroxymethylfurfural in hot compressed water by microwave heating, *Catalysis Communications*, 9 (2008) 2244-2249.
- [77] H. Yan, Y. Yang, D. Tong, X. Xiang, C. Hu, Catalytic conversion of glucose to 5-hydroxymethylfurfural over SO₄²⁻/ZrO₂ and SO₄²⁻/ZrO₂ – Al₂O₃ solid acid catalysts, *Catalysis Communications*, 10 (2009) 1558-1563.
- [78] K. Yamaguchi, T. Sakurada, Y. Ogasawara, N. Mizuno, Tin-tungsten mixed oxide as efficient heterogeneous catalyst for conversion of saccharides to furan derivatives, *Chemistry Letters*, 40 (2011) 542-543.
- [79] S. Lima, A.S. Dias, Z. Lin, P. Brandão, P. Ferreira, M. Pillinger, J. Rocha, V. Calvino-Casilda, A.A. Valente, Isomerization of D-glucose to D-fructose over metallosilicate solid bases, *Applied Catalysis A: General*, 339 (2008) 21-27.
- [80] G. Yang, E.A. Pidko, E.J. Hensen, The Mechanism of Glucose Isomerization to Fructose over Sn-BEA Zeolite: A Periodic Density Functional Theory Study, *ChemSusChem*, 6 (2013) 1688-1696.
- [81] N. Rajabbeigi, A.I. Torres, C.M. Lew, B. Elyassi, L. Ren, Z. Wang, H.J. Cho, W. Fan, P. Daoutidis, M. Tsapatsis, On the kinetics of the isomerization of glucose to fructose using Sn-Beta, *Chemical Engineering Science*, 116 (2014) 235-242.
- [82] R. Bermejo-Deval, R.S. Assary, E. Nikolla, M. Moliner, Y. Román-Leshkov, S.-J. Hwang, A. Palsdóttir, D. Silverman, R.F. Lobo, L.A. Curtiss, Metalloenzyme-like catalyzed isomerizations of sugars by Lewis acid zeolites, *Proceedings of the National Academy of Sciences*, 109 (2012) 9727-9732.

- [83] G. Sørensen, K.B. Sørensen, H.O. Hansen, S.D. Nygaard, Fuel for the Future: Development of New Fuels, eg Biofuels, Applied Microbiology and Molecular Biology in Oilfield Systems, Springer, 2011, pp. 219-228.
- [84] J. Sanders, J. Clark, G. Harmsen, H. Heeres, J. Heijnen, S. Kersten, van Swaaij WPM, Moulijn JA (2012) Process intensification in the future production of base chemicals from biomass, Chem Eng Process, 51 (2012) 117-136.
- [85] X. Qi, M. Watanabe, T.M. Aida, R.L. Smith Jr, Selective conversion of D-fructose to 5-hydroxymethylfurfural by ion-exchange resin in acetone/dimethyl sulfoxide solvent mixtures, Industrial & Engineering Chemistry Research, 47 (2008) 9234-9239.
- [86] X. Qi, M. Watanabe, T.M. Aida, R.L. Smith Jr, Catalytic dehydration of fructose into 5-hydroxymethylfurfural by ion-exchange resin in mixed-aqueous system by microwave heating, Green Chemistry, 10 (2008) 799-805.
- [87] V. Ordonsky, J. Van der Schaaf, J. Schouten, T. Nijhuis, The effect of solvent addition on fructose dehydration to 5-hydroxymethylfurfural in biphasic system over zeolites, Journal of Catalysis, 287 (2012) 68-75.
- [88] M.M. Antunes, S. Lima, M. Pillinger, A.A. Valente, Coupling of nanoporous chromium, aluminium-containing silicates with an ionic liquid for the transformation of glucose into 5-(Hydroxymethyl)-2-furaldehyde, Molecules, 17 (2012) 3690-3707.
- [89] Y. Zhang, V. Degirmenci, C. Li, E.J. Hensen, Phosphotungstic Acid Encapsulated in Metal-Organic Framework as Catalysts for Carbohydrate Dehydration to 5-Hydroxymethylfurfural, ChemSusChem, 4 (2011) 59-64.
- [90] N. Testova, E. Paukshtis, K. Ione, Some peculiarities in synthesis of nitrogen-containing heterocyclic compounds on zeolite catalysts, Reaction Kinetics and Catalysis Letters, 44 (1991) 243-249.
- [91] 周硼, 硫酸衍生固体酸-负载型硫酸及其盐和磺酸树脂催化性能的研究, 大连理工大学, 2003.
- [92] J. Zhao, G. Wang, L. Qin, H. Li, Y. Chen, B. Liu, Synthesis and catalytic cracking performance of mesoporous zeolite Y, Catalysis Communications, 73 (2016) 98-102.
- [93] J. Zhao, Y. Yin, Y. Li, W. Chen, B. Liu, Synthesis and characterization of mesoporous zeolite Y by using block copolymers as templates, Chemical Engineering Journal, 284 (2016) 405-411.
- [94] M.J. Antal, T. Leesomboon, W.S. Mok, G.N. Richards, Mechanism of formation of 2-furaldehyde from D-xylose, Carbohydrate Research, 217 (1991) 71-85.
- [95] J.N. Chheda, Y. Román-Leshkov, J.A. Dumesic, Production of 5-hydroxymethylfurfural and furfural by dehydration of biomass-derived mono-and poly-saccharides, Green Chemistry, 9 (2007) 342-350.
- [96] 范芸珠, 曹发海, 硫酸铵热分解反应动力学研究, 高校化学工程学报, 25 (2011) 341-346.
- [97] 庄新姝, 王树荣, 袁振宏, 骆仲决, 吴创之, 岑可法, 纤维素超低酸水解产物的分析, 农业工程学报, 23 (2007) 177-182.
- [98] T.V. Ojumu, E. Betiku, B.O. Solomon, Auto-hydrolysis of lignocellulosics under extremely low sulphuric acid and high temperature conditions in batch reactor, Biotechnology and Bioprocess Engineering, 8 (2003) 291-293.
- [99] J.S. Kim, Y. Lee, R.W. Torget, Cellulose hydrolysis under extremely low sulfuric acid and high-temperature conditions, Applied biochemistry and biotechnology, 91 (2001) 331-340.
- [100] B.Y. Yang, R. Montgomery, Alkaline degradation of glucose: effect of initial concentration of reactants, Carbohydrate Research, 280 (1996) 27-45.
- [101] J. De Bruijn, A. Kieboom, H. Van Bakkum, Alkaline degradation of monosaccharides III. Influence of reaction parameters upon the final product composition, Recueil des Travaux Chimiques des Pays-Bas, 105 (1986) 176-183.
- [102] M. Orozco, F.J. Luque, Theoretical methods for the description of the solvent effect in biomolecular systems, Chemical Reviews, 100 (2000) 4187-4226.
- [103] Y.B. Tewari, Thermodynamics of industrially-important, enzyme-catalyzed reactions, Applied biochemistry and biotechnology, 23 (1990) 187-203.

- [104] E.H. Ajandouz, A. Puigserver. Nonenzymatic browning reaction of essential amino acids: effect of pH on caramelization and Maillard reaction kinetics, *Journal of Agricultural & Food Chemistry*, 47 (1999) 1786-1793.
- [105] D.A. Ledward, *Developments in food proteins-1* Edited by B. J. F. Hudson, Applied Science Publishers Ltd, London. 335 pp. 1982. Price: £ 35•00, *Meat Science*, 7 (1982) 242 – 243.
- [106] S.H. Wettlaufer, A.C. Leopold, Relevance of Amadori and Maillard products to seed deterioration, *Plant Physiology*, 97 (1991) 165-169.
- [107] T. Oliviero, E. Capuano, B. Cämmerer, V. Fogliano, Influence of roasting on the antioxidant activity and HMF formation of a cocoa bean model systems, *Journal of agricultural and food chemistry*, 57 (2008) 147-152.
- [108] D.S. Mottram, B.L. Wedzicha, A.T. Dodson, Acrylamide is formed in the Maillard reaction, *Nature*, 419 (2002) p á gs. 448-449.
- [109] J.M. Ames, A. Apriyantono, A. Arnoldi, Low molecular weight coloured compounds formed in xylose-lysine model systems, *Food Chemistry*, 46 (1993) 121-127.
- [110] H.-L. Lin, W.-H. Liu, Y.-F. Liu, C.-H. Cheng, Complexation equilibrium constants of poly (vinyl alcohol)-borax dilute aqueous solutions-consideration of electrostatic charge repulsion and free ions charge shielding effect, *Journal of Polymer Research*, 9 (2002) 233-238.
- [111] H. Ochiai, S. Shimizu, Y. Tadokoro, I. Murakami, Complex formation between poly (vinyl alcohol) and borate ion, *Polymer*, 22 (1981) 1456-1458.
- [112] H. Kurokawa, M. Shibayama, T. Ishimaru, S. Nomura, W.-I. Wu, Phase behaviour and sol-gel transition of poly (vinyl alcohol)-borate complex in aqueous solution, *Polymer*, 33 (1992) 2182-2188.
- [113] L. Leibler, E. Pezron, P. Pincus, Viscosity behaviour of polymer solutions in the presence of complexing ions, *Polymer*, 29 (1988) 1105-1109.
- [114] G. Keita, A. Ricard, R. Audebert, E. Pezron, L. Leibler, The poly (vinyl alcohol)-borate system: influence of polyelectrolyte effects on phase diagrams, *Polymer*, 36 (1995) 49-54.
- [115] F. Ilgen, D. Ott, D. Kralisch, C. Reil, A. Palmberger, B. König, Conversion of carbohydrates into 5-hydroxymethylfurfural in highly concentrated low melting mixtures, *Green Chemistry*, 11 (2009) 1948-1954.
- [116] F. Liu, J. Barrault, K. De Oliveira Vigier, F. J é rôme, Dehydration of Highly Concentrated Solutions of Fructose to 5-Hydroxymethylfurfural in a Cheap and Sustainable Choline Chloride/Carbon Dioxide System, *ChemSusChem*, 5 (2012) 1223-1226.
- [117] F. Liu, M. Audemar, K.D.O. Vigier, D. Cartigny, J.-M. Clacens, M.F.C. Gomes, A.A. P á dua, F. De Campo, F. J é rôme, Selectivity enhancement in the aqueous acid-catalyzed conversion of glucose to 5-hydroxymethylfurfural induced by choline chloride, *Green Chemistry*, 15 (2013) 3205-3213.
- [118] Q. Zhang, K.D.O. Vigier, S. Royer, F. J é rôme, Deep eutectic solvents: syntheses, properties and applications, *Chemical Society Reviews*, 41 (2012) 7108-7146.
- [119] L. Hu, Y. Sun, L. Lin, Efficient conversion of glucose into 5-hydroxymethylfurfural by chromium (III) chloride in inexpensive ionic liquid, *Industrial & Engineering Chemistry Research*, 51 (2012) 1099-1104.
- [120] T. Ståhlberg, S. Rodriguez-Rodriguez, P. Fristrup, A. Riisager, Metal-Free Dehydration of Glucose to 5-(Hydroxymethyl) furfural in Ionic Liquids with Boric Acid as a Promoter, *Chemistry-A European Journal*, 17 (2011) 1456-1464.
- [121] M. Francisco, A. van den Bruinhorst, M.C. Kroon, Low-transition-temperature mixtures (LTTMs): A new generation of designer solvents, *Angewandte Chemie International Edition*, 52 (2013) 3074-3085.

Degree papers are in the “[Xiamen University Electronic Theses and Dissertations Database](#)”.

Fulltexts are available in the following ways:

1. If your library is a CALIS member libraries, please log on <http://etd.calis.edu.cn/> and submit requests online, or consult the interlibrary loan department in your library.
2. For users of non-CALIS member libraries, please mail to etd@xmu.edu.cn for delivery details.